

## Evidence-Based Treatment of Tennis Elbow

Smidt N, van der Windt DA, Assendelft WJ, et al. Corticosteroid injections, physiotherapy, or a wait-and-see policy for lateral epicondylitis: a randomised controlled trial. *Lancet* 2002;359:657–62.

### Study Overview

**Objectives.** To compare 3 common treatments for lateral epicondylitis and determine which is the most effective.

**Design.** Randomized controlled trial with an intention-to-treat analysis.

**Setting and participants.** Consecutive participants were recruited by 85 family doctors and referred to 1 of 5 research centers in the Netherlands. Inclusion criteria included pain at the lateral side of the elbow; increasing pain with pressure on the lateral epicondyle and resisted dorsiflexion of the wrist; age 18 to 70 years; ability to complete the questionnaires; and willingness to participate. Exclusion criteria included treatment of elbow complaints with corticosteroid injections or physiotherapy within the preceding 6 months; bilateral elbow symptoms, duration of pain less than 6 weeks, signs and symptoms to suggest another cause of elbow pain, prior surgery of the elbow, dislocation, tendon ruptures, or fractures in the area over the prior year; systemic musculoskeletal or neurological disorders; and contraindications to corticosteroids.

**Intervention.** The intervention occurred over a 6-week period and compared corticosteroid injections and physiotherapy to a control treatment of wait-and-see. Patients assigned to injections received 1 mL (10 mg/mL) of triamcinolone acetone and 1 mL lidocaine 2%. Patients were injected at every tender spot within the elbow until the patient was free of pain during resisted dorsiflexion. During the 6-week intervention period, a maximum of 3 injections was recommended. All providers received training in injection technique prior to the trial. Patients assigned to physiotherapy underwent 9 treatments of pulsed ultrasound, deep friction massage, and an exercise program over the 6-week intervention period. All physiotherapists received special training in the intervention techniques. Wait-and-see patients were treated with analgesic medications and encouragement.

**Main outcome measures.** Outcomes were assessed before randomization and at 3, 6, 12, 26, and 52 weeks after randomization. Most primary outcomes were self-reported by

the patient and assessed through questionnaires. These outcomes included pain during the day, severity of the main complaint, inconvenience, elbow disability, and patient satisfaction. The primary outcome of overall severity of elbow complaints was scored by a blinded research physiotherapist after taking a standardized history and physical examination. Secondary outcomes included pain-free grip strength, maximal grip strength, and pressure pain threshold, and they were measured by a research physiotherapist using standardized equipment.

**Main results.** 185/259 patients referred to the research center were randomized. The dropout rate after randomization was less than 2%. The groups were similar, except that participants randomized to steroid injections had more previous episodes of lateral elbow pain and more concomitant neck disorders. At 6 weeks, 57 patients (92%) in the injection group, 30 (47%) in the physiotherapy group, and 19 (32%) in the wait-and-see group noted significant improvement in their symptoms. Additionally, the corticosteroid injection group scored significantly better in all primary and secondary outcomes compared with physiotherapy and wait-and-see. However, by 26 and 52 weeks, the physiotherapy group scored significantly better than the injection therapy group in all outcomes. Physiotherapy was slightly superior to wait-and-see, but the difference was nonsignificant.

**Conclusion.** For lateral epicondylitis, corticosteroid injections seem to offer immediate and short-term improvement, but more sustained improvement results from physiotherapy.

### Commentary

Lateral epicondylitis is a common complaint affecting between 1% and 2% of the general population [1]. Despite this frequency, very little evidence exists to guide clinicians. A meta-analysis on treatment for lateral epicondylitis concluded

*"Outcomes Research in Review" is edited by Harvey J. Murff, MD, Stephen D. Persell, MD, Josh F. Peterson, MD, and Eric G. Poon, MD, all from the Department of Medicine, Brigham and Women's Hospital; and by David R. Spigel, MD, from the Dana Farber Cancer Institute, Boston, MA.*

that there was insufficient evidence to support any of the currently utilized therapies for lateral epicondylitis [2]. Still, the practitioners are left with several potential therapies for tennis elbow but no good evidence to direct them to the best practice. Therefore, Smidt et al's well-designed randomized controlled study is a welcomed contribution to the literature. Despite the difficulties in blinding the research physiotherapist to the treatment allocations, numerous carefully constructed methodologic safeguards were implemented to minimize biases.

However, there are still flaws in the study. Enrollments into the study were based on physician selection and were not complete during the study process. According to participating physicians, most eligible patients were missed because of provider forgetfulness or busy office hours; it is unlikely that this would influence the overall results. Also, despite attempts to keep the research physiotherapists blinded, they correctly guessed treatment allocations in 52% of the patients. The authors note that the physiotherapists generally based their guesses concerning treatment allocation on the results of the outcome assessment. If this were the case, then

it is less likely that this partial unblinding would influence the results because the outcome would have been measured prior to discovery of the treatment allocation.

### Applications for Clinical Practice

When discussing treatment options for lateral epicondylitis, it is important to inform patients that different treatment strategies confer a different duration of benefit. Corticosteroid injections seem to offer more short-term benefit that disappears over time, while the benefits of physiotherapy become more evident after several months.

*—Review by Harvey J. Murff, MD*

### References

1. Verhaar JA. Tennis elbow. Anatomical, epidemiological and therapeutic aspects. *Int Orthop* 1994;18:263–7.
2. Labelle H, Guibert R, Joncas J, et al. Lack of scientific evidence for the treatment of lateral epicondylitis of the elbow. An attempted meta-analysis. *J Bone Joint Surg Br* 1992;74:646–51.

Copyright 2002 by Turner White Communications Inc., Wayne, PA. All rights reserved.