Key to abbreviations: *abs/comm* = abstract with commentary.

SUBJECTS

Abdominal pain
Functional abdominal pain in children, 15(5): 248–58
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Acne
Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*
Acute coronary syndrome
Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*
Adapalene
Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*
Alzheimer's disease
Diagnosis and treatment of Alzheimer's disease, 15(2):87–100
Anorexia nervosa
Anorexia nervosa: role of the primary care physician, 15(9):450–60
Anticoagulation
Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90
Anxiety
Generalized anxiety disorder in adults, 15(7): 353–63
Aortic stenosis
Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
Aspirin
No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575–6 *abs/comm*
Asthma
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Barrett's esophagus
Barrett's esophagus: evaluation and management, 15(8):402–11
Behavioral therapy
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
Benzoyl peroxide
Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*
Beta blockers
The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10): 470–1 *abs/comm*
Beta carotene
Beta carotene supplementation may improve cognitive function over the long term, 15(1): 10–1 *abs/comm*
Biologic therapy
Biologic agents in the treatment of Crohn's disease, 15(1):31–8
Biomarkers
Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112–3 *abs/comm*
Bivalirudin
Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*
Blood glucose
Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7): 318–9 *abs/comm*
Cancer
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6): 272–4 *abs/comm*
End-of-life discussions associated with less aggressive care in patients dying of cancer, 15(12):583–5
Prostate cancer screening: benefits remain uncertain, 15(10):474–5 *abs/comm*
Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*
Stool DNA testing for colorectal cancer shows promise, 15(12):578–9 *abs/comm*
A young woman with a thyroid nodule, 15(9): 443–9
Cardiac arrest
Chest compression–only CPR as effective as standard CPR, 15(2):60–1 *abs/comm*
CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371–2 *abs/comm*
Cardiovascular disease
Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112–3 *abs/comm*
Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531–2 *abs/comm*
Cardiovascular risk
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46
Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5):220–1 *abs/comm*
Case management
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Catheterization
Femoral versus jugular central venous line placement: which is better?, 15(8):378–9 *abs/comm*
Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*
Certification examination
Do higher tests scores equate to higher-quality care?, 15(9):426–7 *abs/comm*
Cirrhosis
Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400
Primary care management of the patient with cirrhosis, 15(11):554–63
Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*
Clinical pathway
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
Coffee
Another reason to drink your morning cup of coffee, 15(9):429–30 *abs/comm*
Cognitive function
Beta carotene supplementation may improve cognitive function over the long term, 15(1): 10–1 *abs/comm*
Collaborative research
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Computed tomography
Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*
Coronary artery disease
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52
Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5): 220–1 *abs/comm*
Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*
Corticosteroids
Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268–9 *abs/comm*
CPR
Chest compression–only CPR as effective as standard CPR, 15(2):60–1 *abs/comm*

CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371–2 *abs/comm*
Crohn's disease
Biologic agents in the treatment of Crohn's disease, 15(1):31–8
Deep venous thrombosis
Link between air pollution and deep venous thrombosis risk, 15(7):326–7 *abs/comm*
Delirium
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502–11
Dementia
Diagnosis and treatment of Alzheimer's disease, 15(2):87–100
Diabetes
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Diabetes group education and self-management intervention shows mixed results, 15(5):225–6 *abs/comm*
Early intervention to prevent type 2 diabetes, 15(1):39–47
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12): 575–6 *abs/comm*
Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7): 318–9 *abs/comm*
Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427–9 *abs/comm*
Diet
Another reason to drink your morning cup of coffee, 15(9):429–30 *abs/comm*
Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427–9 *abs/comm*
DNA testing
Stool DNA testing for colorectal cancer shows promise, 15(12):578–9 *abs/comm*
Drug-eluting stents
No differences found between drug-eluting stents in everyday clinical practice, 15(3): 113–4 *abs/comm*
Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*
Drug review
FDA drug review deadlines: a safety concern?, 15(5):224–5 *abs/comm*
Dyslipidemia
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*
Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323–4 *abs/comm*
Eating disorders
Anorexia nervosa: role of the primary care physician, 15(9):450–60
Electronic health record
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Ezetimibe
Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
Falls
Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Fracture
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94
Gastrointestinal disorders
Functional abdominal pain in children, 15(5): 248–58
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Generalized anxiety disorder
Generalized anxiety disorder in adults, 15(7): 353–63
GERD
Barrett's esophagus: evaluation and management, 15(8):402–11
Geriatric care
Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
Low-income seniors may benefit from a home-based primary care program, 15(2):64–5 *abs/comm*
Glucosamine
Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219–20 *abs/comm*
Glycemic control
Early intervention to prevent type 2 diabetes, 15(1):39–47
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Graves' disease
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
Guidelines
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Headache
Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*
Health disparities
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Hearing impairment
Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
Heart failure
N-3 polyunsaturated fatty acids reduce heart failure–related admissions and death, 15(11): 522–3 *abs/comm*
Hepatitis B
Evaluation and management of hepatitis B virus infection, 15(3):147–55
Home care
Low-income seniors may benefit from a home-based primary care program, 15(2):64–5 *abs/comm*
Hospital performance
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Hospitalist
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Hydrocortisone
No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116–7 *abs/comm*
Hypertension
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
Hyperthyroidism
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
Influenza
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Insomnia
Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203
Insurance
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6): 272–4 *abs/comm*
Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*
Lipid lowering
Limited benefit for aggressive lipid lowering

with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
Liver disease
Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400
Primary care management of the patient with cirrhosis, 15(11):554–63
Lumbar puncture
Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*
Measurement error
Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323–4 *abs/comm*
Medicaid
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272–4 *abs/comm*
Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*
Medical education
A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30
Medical error
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
Medication adherence
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Meningitis
Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*
Meta-analysis
Meta-analyses: caveat lector, 15(2):61–2 *abs/comm*
Migraine
Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*
Myocardial infarction
Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*
Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*
Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*
Noncardiac surgery
The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470–1 *abs/comm*
Nosocomial infections
Femoral versus jugular central venous line placement: which is better?, 15(8):378–9 *abs/comm*
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85
Obesity
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
Omega-3 fatty acids
N-3 polyunsaturated fatty acids reduce heart failure–related admissions and death, 15(11):522–3 *abs/comm*
Osteoarthritis
Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219–20 *abs/comm*
Osteoporosis
Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94
Patent foramen ovale
Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*
Patient-centered care
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Patient safety
FDA drug review deadlines: a safety concern?, 15(5):224–5 *abs/comm*
To err is human continued: a failure of follow-up, 15(1):21–3
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Percutaneous coronary intervention
Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52
Emergency physician–initiated percutaneous coronary intervention and effects on door-to-balloon time, 15(2):63–4 *abs/comm*
No differences found between drug-eluting stents in everyday clinical practice, 15(3):113–4 *abs/comm*
Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*
Perioperative care
The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470–1 *abs/comm*
Peripheral arterial disease
No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575–6 *abs/comm*
Pharmacogenetics
Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90
Pneumococcal vaccine
Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*
Pneumonia
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85
Pollution
Link between air pollution and deep venous thrombosis risk, 15(7):326–7 *abs/comm*
Prediabetes
Early intervention to prevent type 2 diabetes, 15(1):39–47
Preeclampsia
Diagnosis and management of preeclampsia, 15(12):607–14
Provider support
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
Quality of care
Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*
Reperfusion therapy
Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*
Reports from the Field
A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Resource utilization
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Revascularization
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52
Rheumatoid arthritis
Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268–9 *abs/comm*
Risk assessment
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*
Screening
Prostate cancer screening: benefits remain uncertain, 15(10):474–5 *abs/comm*
Self-management
Diabetes group education and self-management intervention shows mixed results, 15(5):225–6 *abs/comm*
Septic shock
No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116–7 *abs/comm*
Sleep disorders
Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203
Smoking cessation
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92

From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
New and emerging treatments in smoking cessation, 15(11):545–52
Stents
Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*
No differences found between drug-eluting stents in everyday clinical practice, 15(3):113–4 *abs/comm*
Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*
Subarachnoid hemorrhage
Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*
Supplementation
Beta carotene supplementation may improve cognitive function over the long term, 15(1):10–1 *abs/comm*
Telemetry
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Test results management
To err is human continued: a failure of follow-up, 15(1):21–3
Thiazolidinediones
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Thyroid nodule
A young woman with a thyroid nodule, 15(9):443–9
Thyroiditis
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
Tobacco use
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
New and emerging treatments in smoking cessation, 15(11):545–52
Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*
Underserved populations
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Vaccination
Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Vitamin D
Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*
Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531–2 *abs/comm*
Weight loss
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44

AUTHORS

Abad C
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85
Abudayyeh S
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Adamson SC
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66–70
An LC
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Anandasabapathy S
Barrett's esophagus: evaluation and management, 15(8):402–11
Anderson DC Jr
Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90
Anton SD
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
Arcury TA
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Arnedt T
Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203
Attia E
Anorexia nervosa: role of the primary care physician, 15(9):450–60
Barbieri KE
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Bartel GA
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66–70
Bernardo LM
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Bernhardt T
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Bertoni AG
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Bitton A
Diabetes group education and self-management intervention shows mixed results, 15(5):225–6 *abs/comm*
Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*
N-3 polyunsaturated fatty acids reduce heart failure–related admissions and death, 15(11):522–3 *abs/comm*
Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268–9 *abs/comm*
Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7):318–9 *abs/comm*
Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*
Stool DNA testing for colorectal cancer shows promise, 15(12):578–9 *abs/comm*
Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427–9 *abs/comm*
Block JP
Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*
End-of-life discussions associated with less aggressive care in patients dying of cancer, 15(12):583–5
Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323–4 *abs/comm*
Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219–20 *abs/comm*
Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*
Prostate cancer screening: benefits remain uncertain, 15(10):474–5 *abs/comm*
Bluhm J
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Boyle DJ
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
Brown SJ
Biologic agents in the treatment of Crohn's disease, 15(1):31–8
Bulger J
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Callanan MG
To err is human continued: a failure of follow-up, 15(1):21–3
Carrier ER
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272–4 *abs/comm*
Carty M
A program for reducing noneducational resident

activities at an academic medical center, 15(1): 25–30
Cavanaugh J
To err is human continued: a failure of follow-up, 15(1):21–3
Center B
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Chin MH
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Chung JA
Diagnosis and treatment of Alzheimer's disease, 15(2):87–100
Cogen FR
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
Cotler SJ
Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400
Dedhia P
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
DeGuzman PB
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Diduszyn J
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Dolor RJ
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Drum ML
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Edwards-Lee T
Diagnosis and treatment of Alzheimer's disease, 15(2):87–100
Engberg S
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Fares R
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Ferrari P
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Ferretti J
Isopropyl alcohol left on the skin falsely lowers capillary glucose values, 15(4):179–81
Finn KM
A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30
Finstad D
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Foldes S
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Fraley JK
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Fremont AM
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Friedberg MW
Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371–2 *abs/comm*
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
Low-income seniors may benefit from a home-based primary care program, 15(2):64–5 *abs/comm*
No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12): 575–6 *abs/comm*
No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116–7 *abs/comm*
Nurse-led counseling had no effect on heart failure outcomes, 15(4):170–1 *abs/comm*
Furst J
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66–70
Gallagher TH
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
Gilger MA
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Goldberg KC
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Graham DY
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Grice GR
Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90
Grzywacz JG
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Gurm HS
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52
Hafner JM
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Harrington MB
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502–11
Harris L
A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30
Harrison JF
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Hellkamp AS
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Hess R
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Hinson T
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Hofmann MT
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Howell E
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Huang RL
Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*
Chest compression–only CPR as effective as standard CPR, 15(2):60–1 *abs/comm*
Ezetimibe's clinical benefit called into question, 15(6):277–8 *abs/comm*
No differences found between drug-eluting stents in everyday clinical practice, 15(3): 113–4 *abs/comm*
Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*
Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*
Hud-Aleem R
Generalized anxiety disorder in adults, 15(7): 353–63
Hwang U
Emergency physician–initiated percutaneous coronary intervention and effects on door-to-balloon time, 15(2):63–4 *abs/comm*
Femoral versus jugular central venous line placement: which is better?, 15(8):378–9 *abs/comm*
Racial and ethnic disparities continue for pain care in U.S. emergency departments, 15(4): 173–4 *abs/comm*
Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*
Hyams JS
Functional abdominal pain in children, 15(5): 248–58
Ingram KD
Primary care management of the patient with cirrhosis, 15(11):554–63
Jain SH
Early intervention to prevent type 2 diabetes, 15(1):39–47

Jha AK
Beta carotene supplementation may improve cognitive function over the long term, 15(1): 10–1 *abs/comm*
Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112–3 *abs/comm*
Do higher tests scores equate to higher-quality care?, 15(9):426–7 *abs/comm*
Link between air pollution and deep venous thrombosis risk, 15(7):326–7 *abs/comm*
Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531–2 *abs/comm*
Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5):220–1 *abs/comm*
Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*
Johnson PA
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
Jones JB
Meta-analyses: caveat lector, 15(2):61–2 *abs/comm*
Julien S
Diagnosis and management of preeclampsia, 15(12):607–14
Kallwitz ER
Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400
Kamat PP
Barrett's esophagus: evaluation and management, 15(8):402–11
Kanwar M
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Kaplowitz PB
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
Katz JT
A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30
Keyhani S
Another reason to drink your morning cup of coffee, 15(9):429–30 *abs/comm*
Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*
FDA drug review deadlines: a safety concern?, 15(5):224–5 *abs/comm*
Kirch MA
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Kirchhoff AC
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Kirk JK
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Klem ML
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Koss RG
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Kravet S
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
Kumar CB
Intensive intervention reduces mortality in type 2 diabetes, 15(4):171–3 *abs/comm*
Ladson D
Generalized anxiety disorder in adults, 15(7): 353–63
Laing M
A program for reducing noneducational resident activities at an academic medical center, 15(1): 25–30
Lamb P
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Laramie D
Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90
Lee JC
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Levie J
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Lippold S
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Litvin C
To err is human continued: a failure of follow-up, 15(1):21–3
Loeb JM
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Malaty HM
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Manley M
Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92
Martin KD
Isopropyl alcohol left on the skin falsely lowers capillary glucose values, 15(4):179–81
McCotter P
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
McKinley L
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85
McMahon GT
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46
Megginson GC
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Milsom VA
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
Minnick S
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Montgomery D
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Morton DJ
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Mui C
New and emerging treatments in smoking cessation, 15(11):545–52
The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470–1 *abs/comm*
Murff HJ
Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1): 9–10 *abs/comm*
Musante D
Anorexia nervosa: role of the primary care physician, 15(9):450–60
Nackers LM
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
Naglak M
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Newton RL Jr
Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44
O'Malley K
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Organist LA
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Pantoja PM
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Parks CM
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Pastor W
Length of stay safely reduced with a diabetes

clinical pathway in an urban children's hospital, 15(8):384–92
Patterson B
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Pratt DS
Evaluation and management of hepatitis B virus infection, 15(3):147–55
Rachelefsky L
A program for reducing noneducational resident activities at an academic medical center, 15(1): 25–30
Raji A
Early intervention to prevent type 2 diabetes, 15(1):39–47
Rand E
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Reed MC
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52
Rohrer JE
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66–70
Rosman HS
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Rubal BJ
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Rudolph JL
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502–11
Ryan C
Optimizing telemetry utilization in an academic medical center, 15(9):435–40
Safdar N
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2): 71–85
Saravolatz L
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16–20
Schaefer CT
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Schafer AL
Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94
Schlenk EA
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Schlichting JA
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Schreiber K
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502–11
Sekhobo JP
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Sellmeyer DE
Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94
Setodji C
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Shah NR
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6): 272–4 *abs/comm*
Intensive intervention reduces mortality in type 2 diabetes, 15(4):171–3 *abs/comm*
Meta-analyses: caveat lector, 15(2):61–2 *abs/comm*
The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10): 470–1 *abs/comm*
Sherman SE
New and emerging treatments in smoking cessation, 15(11):545–52
Simpson AG
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Siraj ES
A young woman with a thyroid nodule, 15(9): 443–9
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
Smith A
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Smith DG
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
Sun BC
Patient-centered care and outcomes after myocardial infarction, 15(3):119–24
Swanson LM
Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203
Tenner CT
To err is human continued: a failure of follow-up, 15(1):21–3
Tkachenko M
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
Trikudanathan S
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46
Triola MM
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*
Veloski C
A young woman with a thyroid nodule, 15(9): 443–9
Viette CR
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
Wade T
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66–70
Walter EB
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
Wang C
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Waterman AD
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
White AA
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7
Williams SC
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
Wilson J
Biologic agents in the treatment of Crohn's disease, 15(1):31–8
Wright S
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
You WB
Diagnosis and management of preeclampsia, 15(12):607–14
Youssef NN
Functional abdominal pain in children, 15(5): 248–58
Zaman
Primary care management of the patient with cirrhosis, 15(11):554–63
Zhang JX
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31

Copyright 2008 by Turner White Communications Inc., Wayne, PA. All rights reserved.