

Key to abbreviations: *abs/comm* = abstract with commentary.

SUBJECTS

Abdominal pain

Functional abdominal pain in children, 15(5): 248–58

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92

Acne

Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*

Acute coronary syndrome

Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*

Adapalene

Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*

Alzheimer's disease

Diagnosis and treatment of Alzheimer's disease, 15(2):87–100

Anorexia nervosa

Anorexia nervosa: role of the primary care physician, 15(9):450–60

Anticoagulation

Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90

Anxiety

Generalized anxiety disorder in adults, 15(7): 353–63

Aortic stenosis

Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*

Aspirin

No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575–6 *abs/comm*

Asthma

Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34

Barrett's esophagus

Barrett's esophagus: evaluation and management, 15(8):402–11

Behavioral therapy

Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44

Benzoyl peroxide

Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9–10 *abs/comm*

Beta blockers

The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10): 470–1 *abs/comm*

Beta carotene

Beta carotene supplementation may improve cognitive function over the long term, 15(1): 10–1 *abs/comm*

Biologic therapy

Biologic agents in the treatment of Crohn's disease, 15(1):31–8

Biomarkers

Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112–3 *abs/comm*

Bivalirudin

Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*

Blood glucose

Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7): 318–9 *abs/comm*

Cancer

Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6): 272–4 *abs/comm*

End-of-life discussions associated with less aggressive care in patients dying of cancer, 15(12):583–5

Prostate cancer screening: benefits remain uncertain, 15(10):474–5 *abs/comm*

Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*

Stool DNA testing for colorectal cancer shows promise, 15(12):578–9 *abs/comm*

A young woman with a thyroid nodule, 15(9): 443–9

Cardiac arrest

Chest compression–only CPR as effective as standard CPR, 15(2):60–1 *abs/comm*

CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371–2 *abs/comm*

Cardiovascular disease

Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112–3 *abs/comm*

Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531–2 *abs/comm*

Cardiovascular risk

Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*

Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46

Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5):220–1 *abs/comm*

Case management

Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501

Catheterization

Femoral versus jugular central venous line placement: which is better?, 15(8):378–9 *abs/comm*

Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*

Certification examination

Do higher tests scores equate to higher-quality care?, 15(9):426–7 *abs/comm*

Cirrhosis

Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400

Primary care management of the patient with cirrhosis, 15(11):554–63

Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*

Clinical pathway

Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92

Coffee

Another reason to drink your morning cup of coffee, 15(9):429–30 *abs/comm*

Cognitive function

Beta carotene supplementation may improve cognitive function over the long term, 15(1): 10–1 *abs/comm*

Collaborative research

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9

Computed tomography

Revisiting the need for computed tomography / lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*

Coronary artery disease

Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52

Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*

Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43

Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5): 220–1 *abs/comm*

Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*

Corticosteroids

Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268–9 *abs/comm*

CPR

Chest compression–only CPR as effective as standard CPR, 15(2):60–1 *abs/comm*

- CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371–2 *abs/comm*
- Crohn's disease**
Biologic agents in the treatment of Crohn's disease, 15(1):31–8
- Deep venous thrombosis**
Link between air pollution and deep venous thrombosis risk, 15(7):326–7 *abs/comm*
- Delirium**
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502–11
- Dementia**
Diagnosis and treatment of Alzheimer's disease, 15(2):87–100
- Diabetes**
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133–46
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Diabetes group education and self-management intervention shows mixed results, 15(5):225–6 *abs/comm*
Early intervention to prevent type 2 diabetes, 15(1):39–47
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281–6
Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384–92
No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575–6 *abs/comm*
Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7):318–9 *abs/comm*
Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427–9 *abs/comm*
- Diet**
Another reason to drink your morning cup of coffee, 15(9):429–30 *abs/comm*
Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427–9 *abs/comm*
- DNA testing**
Stool DNA testing for colorectal cancer shows promise, 15(12):578–9 *abs/comm*
- Drug-eluting stents**
No differences found between drug-eluting stents in everyday clinical practice, 15(3):113–4 *abs/comm*
Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12–3 *abs/comm*
- Drug review**
FDA drug review deadlines: a safety concern?, 15(5):224–5 *abs/comm*
- Dyslipidemia**
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*
Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323–4 *abs/comm*
- Eating disorders**
Anorexia nervosa: role of the primary care physician, 15(9):450–60
- Electronic health record**
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43
- Ezetimibe**
Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
- Falls**
Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6
- Fracture**
Chronic use of thiazolidinediones increases fracture risk, 15(7):322–3 *abs/comm*
Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94
- Gastrointestinal disorders**
Functional abdominal pain in children, 15(5):248–58
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287–92
- Generalized anxiety disorder**
Generalized anxiety disorder in adults, 15(7):353–63
- GERD**
Barrett's esophagus: evaluation and management, 15(8):402–11
- Geriatric care**
Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
Optimizing medication adherence in older patients: a systematic review, 15(12):595–606
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
Low-income seniors may benefit from a home-based primary care program, 15(2):64–5 *abs/comm*
- Glucosamine**
Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219–20 *abs/comm*
- Glycemic control**
Early intervention to prevent type 2 diabetes, 15(1):39–47
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
- Graves' disease**
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
- Guidelines**
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
- Headache**
Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*
- Health disparities**
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494–501
- Hearing impairment**
Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
- Heart failure**
N-3 polyunsaturated fatty acids reduce heart failure-related admissions and death, 15(11):522–3 *abs/comm*
- Hepatitis B**
Evaluation and management of hepatitis B virus infection, 15(3):147–55
- Home care**
Low-income seniors may benefit from a home-based primary care program, 15(2):64–5 *abs/comm*
- Hospital performance**
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329–37
- Hospitalist**
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9
- Hydrocortisone**
No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116–7 *abs/comm*
- Hypertension**
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125–31
Lowering high blood pressure is beneficial in the very elderly, 15(6):276–7 *abs/comm*
- Hyperthyroidism**
Update on the diagnosis and treatment of hyperthyroidism, 15(6):298–307
- Influenza**
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227–34
- Insomnia**
Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203
- Insurance**
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272–4 *abs/comm*
Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*
- Lipid lowering**
Limited benefit for aggressive lipid lowering

with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*

Liver disease

Management of the cirrhotic patient prior to liver transplantation, 15(8):393–400

Primary care management of the patient with cirrhosis, 15(11):554–63

Lumbar puncture

Revisiting the need for computed tomography / lumbar puncture in subarachnoid hemorrhage detection, 15(12):582–3 *abs/comm*

Measurement error

Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323–4 *abs/comm*

Medicaid

Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272–4 *abs/comm*

Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*

Medical education

A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30

Medical error

Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7

Medication adherence

Optimizing medication adherence in older patients: a systematic review, 15(12):595–606

Meningitis

Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*

Meta-analysis

Meta-analyses: caveat lector, 15(2):61–2 *abs/comm*

Migraine

Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*

Myocardial infarction

Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*

Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379–81 *abs/comm*

Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*

Patient-centered care and outcomes after myocardial infarction, 15(3):119–24

Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*

Noncardiac surgery

The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470–1 *abs/comm*

Nosocomial infections

Femoral versus jugular central venous line placement: which is better?, 15(8):378–9 *abs/comm*

Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85

Obesity

Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536–44

Omega-3 fatty acids

N-3 polyunsaturated fatty acids reduce heart failure–related admissions and death, 15(11):522–3 *abs/comm*

Osteoarthritis

Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219–20 *abs/comm*

Osteoporosis

Interventions to improve treatment of osteoporosis following fracture, 15(12):587–94

Patent foramen ovale

Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274–5 *abs/comm*

Patient-centered care

Patient-centered care and outcomes after myocardial infarction, 15(3):119–24

Patient safety

FDA drug review deadlines: a safety concern?, 15(5):224–5 *abs/comm*

To err is human continued: a failure of follow-up, 15(1):21–3

Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6

Percutaneous coronary intervention

Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317–8 *abs/comm*

Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52

Emergency physician–initiated percutaneous coronary intervention and effects on door-to-balloon time, 15(2):63–4 *abs/comm*

No differences found between drug-eluting stents in everyday clinical practice, 15(3):113–4 *abs/comm*

Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*

Perioperative care

The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470–1 *abs/comm*

Peripheral arterial disease

No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575–6 *abs/comm*

Pharmacogenetics

Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183–90

Pneumococcal vaccine

Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373–4 *abs/comm*

Pneumonia

Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71–85

Pollution

Link between air pollution and deep venous thrombosis risk, 15(7):326–7 *abs/comm*

Prediabetes

Early intervention to prevent type 2 diabetes, 15(1):39–47

Preeclampsia

Diagnosis and management of preeclampsia, 15(12):607–14

Provider support

Supporting health care workers after medical error: considerations for health care leaders, 15(5):240–7

Quality of care

Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8–9 *abs/comm*

Reperfusion therapy

Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*

Reports from the Field

A program for reducing noneducational resident activities at an academic medical center, 15(1):25–30

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235–9

Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338–43

Length of stay safely reduced with a diabetes clinical pathway in an urban children’s hospital, 15(8):384–92

Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293–6

Resource utilization

Optimizing telemetry utilization in an academic medical center, 15(9):435–40

Revascularization

Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344–52

Rheumatoid arthritis

Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268–9 *abs/comm*

Risk assessment

Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115–6 *abs/comm*

Screening

Prostate cancer screening: benefits remain uncertain, 15(10):474–5 *abs/comm*

Self-management

Diabetes group education and self-management intervention shows mixed results, 15(5):225–6 *abs/comm*

Septic shock

No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116–7 *abs/comm*

Sleep disorders

Helping patients with insomnia: a guide for the primary care physician, 15(4):191–203

Smoking cessation

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485–92

- From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37
- New and emerging treatments in smoking cessation, 15(11):545-52
- Stents**
- Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*
- No differences found between drug-eluting stents in everyday clinical practice, 15(3):113-4 *abs/comm*
- Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12-3 *abs/comm*
- Subarachnoid hemorrhage**
- Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582-3 *abs/comm*
- Supplementation**
- Beta carotene supplementation may improve cognitive function over the long term, 15(1):10-1 *abs/comm*
- Telemetry**
- Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20
- Optimizing telemetry utilization in an academic medical center, 15(9):435-40
- Test results management**
- To err is human continued: a failure of follow-up, 15(1):21-3
- Thiazolidinediones**
- Chronic use of thiazolidinediones increases fracture risk, 15(7):322-3 *abs/comm*
- Thyroid nodule**
- A young woman with a thyroid nodule, 15(9):443-9
- Thyroiditis**
- Update on the diagnosis and treatment of hyperthyroidism, 15(6):298-307
- Tobacco use**
- Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92
- From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37
- New and emerging treatments in smoking cessation, 15(11):545-52
- Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*
- Underserved populations**
- Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6
- Vaccination**
- Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373-4 *abs/comm*
- Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34
- Vitamin D**
- Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379-81 *abs/comm*
- Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531-2 *abs/comm*
- Weight loss**
- Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536-44
- AUTHORS**
- Abad C**
- Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71-85
- Abudayyeh S**
- Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92
- Adamson SC**
- Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66-70
- An LC**
- Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92
- Anandasabapathy S**
- Barrett's esophagus: evaluation and management, 15(8):402-11
- Anderson DC Jr**
- Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183-90
- Anton SD**
- Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536-44
- Arcury IA**
- Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6
- Ardedt T**
- Helping patients with insomnia: a guide for the primary care physician, 15(4):191-203
- Attia E**
- Anorexia nervosa: role of the primary care physician, 15(9):450-60
- Barbieri KE**
- Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43
- Bartel GA**
- Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66-70
- Bernardo LM**
- Optimizing medication adherence in older patients: a systematic review, 15(12):595-606
- Bernhardt T**
- Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92
- Bertoni AG**
- Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6
- Bitton A**
- Diabetes group education and self-management intervention shows mixed results, 15(5):225-6 *abs/comm*
- Hospitalizations and mortality greatly reduced following introduction of the pneumococcal conjugate vaccine, 15(8):373-4 *abs/comm*
- N-3 polyunsaturated fatty acids reduce heart failure-related admissions and death, 15(11):522-3 *abs/comm*
- Rheumatoid arthritis increases risk of infection requiring hospitalization; risk higher in patients on corticosteroids, 15(6):268-9 *abs/comm*
- Self-monitoring of blood glucose depressingly not helpful in the newly diagnosed, 15(7):318-9 *abs/comm*
- Smoking ban significantly reduces acute coronary syndrome admissions, 15(10):478, 480, 484 *abs/comm*
- Stool DNA testing for colorectal cancer shows promise, 15(12):578-9 *abs/comm*
- Sugar-sweetened beverages increase risk for type 2 diabetes in African-American women, 15(9):427-9 *abs/comm*
- Block JP**
- Deficient 25-hydroxyvitamin D levels are associated with incident cardiovascular events in men, 15(8):379-81 *abs/comm*
- End-of-life discussions associated with less aggressive care in patients dying of cancer, 15(12):583-5
- Frequent cholesterol monitoring prone to error and may be unnecessary, 15(7):323-4 *abs/comm*
- Glucosamine sulfate ineffective for the treatment of hip osteoarthritis, 15(5):219-20 *abs/comm*
- Hearing impairment: another microvascular complication of diabetes?, 15(9):421, 424 *abs/comm*
- Limited benefit for aggressive lipid lowering with simvastatin/ezetimibe in patients with aortic stenosis; possible increased risk of cancer, 15(11):528, 530 *abs/comm*
- Patent foramen ovale closure minimally effective for treating refractory migraines with aura, 15(6):274-5 *abs/comm*
- Prostate cancer screening: benefits remain uncertain, 15(10):474-5 *abs/comm*
- Bluhm J**
- Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92
- Boyle DJ**
- Supporting health care workers after medical error: considerations for health care leaders, 15(5):240-7
- Brown SJ**
- Biologic agents in the treatment of Crohn's disease, 15(1):31-8
- Bulger J**
- Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9
- Callanan MG**
- To err is human continued: a failure of follow-up, 15(1):21-3
- Carrier ER**
- Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272-4 *abs/comm*
- Carty M**
- A program for reducing noneducational resident

activities at an academic medical center, 15(1):25-30

Cavanaugh J

To err is human continued: a failure of follow-up, 15(1):21-3

Center B

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92

Chin MH

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Chung JA

Diagnosis and treatment of Alzheimer's disease, 15(2):87-100

Cogen FR

Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384-92

Cotler SJ

Management of the cirrhotic patient prior to liver transplantation, 15(8):393-400

Dedhia P

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9

DeGuzman PB

Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43

Diduszyn J

Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293-6

Dolor RJ

Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34

Drum ML

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Edwards-Lee T

Diagnosis and treatment of Alzheimer's disease, 15(2):87-100

Engberg S

Optimizing medication adherence in older patients: a systematic review, 15(12):595-606

Fares R

Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20

Ferrari P

Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494-501

Ferretti J

Isopropyl alcohol left on the skin falsely lowers capillary glucose values, 15(4):179-81

Finn KM

A program for reducing noneducational resident activities at an academic medical center, 15(1):25-30

Finstad D

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92

Foldes S

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92

Fraleigh JK

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92

Fremont AM

Patient-centered care and outcomes after myocardial infarction, 15(3):119-24

Friedberg MW

Better use of existing evidence reduces falls among the elderly, 15(10):476, 478 *abs/comm*
CPR was a success, but the patient died: performance debriefing of cardiac arrest processes and postevent outcomes, 15(8):371-2 *abs/comm*

Lowering high blood pressure is beneficial in the very elderly, 15(6):276-7 *abs/comm*

Low-income seniors may benefit from a home-based primary care program, 15(2):64-5 *abs/comm*

No benefit of daily low-dose aspirin in diabetic patients with peripheral arterial disease, 15(12):575-6 *abs/comm*

No mortality benefits with hydrocortisone therapy in patients with septic shock, 15(3):116-7 *abs/comm*

Nurse-led counseling had no effect on heart failure outcomes, 15(4):170-1 *abs/comm*

Furst J

Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66-70

Gallagher TH

Supporting health care workers after medical error: considerations for health care leaders, 15(5):240-7

Gilger MA

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92

Goldberg KC

Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34

Graham DY

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92

Grice GR

Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183-90

Grzywacz JG

Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6

Gurm HS

Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344-52

Hafner JM

From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37

Harrington MB

Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502-11

Harris L

A program for reducing noneducational resident activities at an academic medical center, 15(1):25-30

Harrison JF

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Hellkamp AS

Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34

Hess R

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9

Hinson T

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9

Hofmann MT

Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293-6

Howell E

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9

Huang RL

Bivalirudin reduces adverse events during primary percutaneous coronary intervention for acute myocardial infarction, 15(7):317-8 *abs/comm*

Chest compression-only CPR as effective as standard CPR, 15(2):60-1 *abs/comm*

Ezetimibe's clinical benefit called into question, 15(6):277-8 *abs/comm*

No differences found between drug-eluting stents in everyday clinical practice, 15(3):113-4 *abs/comm*

Real-world use of drug-eluting stents versus bare-metal stents, 15(1):12-3 *abs/comm*

Why is reperfusion therapy withheld in eligible ED patients with STEMI?, 15(5):222 *abs/comm*

Hud-Aleem R

Generalized anxiety disorder in adults, 15(7):353-63

Hwang U

Emergency physician-initiated percutaneous coronary intervention and effects on door-to-balloon time, 15(2):63-4 *abs/comm*

Femoral versus jugular central venous line placement: which is better?, 15(8):378-9 *abs/comm*

Racial and ethnic disparities continue for pain care in U.S. emergency departments, 15(4):173-4 *abs/comm*

Revisiting the need for computed tomography/lumbar puncture in subarachnoid hemorrhage detection, 15(12):582-3 *abs/comm*

Hyams JS

Functional abdominal pain in children, 15(5):248-58

Ingram KD

Primary care management of the patient with cirrhosis, 15(11):554-63

Jain SH

Early intervention to prevent type 2 diabetes, 15(1):39-47

Jha AK

Beta carotene supplementation may improve cognitive function over the long term, 15(1):10-1 *abs/comm*

Can biomarkers improve prediction of cardiovascular events in heart disease patients?, 15(3):112-3 *abs/comm*

Do higher tests scores equate to higher-quality care?, 15(9):426-7 *abs/comm*

Link between air pollution and deep venous thrombosis risk, 15(7):326-7 *abs/comm*

Low levels of 25-hydroxyvitamin D related to increased mortality, but questions remain, 15(11):531-2 *abs/comm*

Near-continuous association between lipoprotein(a) level and CHD risk independent of established cardiovascular risk factors, 15(5):220-1 *abs/comm*

Quality of surveillance for hepatocellular carcinoma determines outcome in cirrhotic patients, 15(4):175 *abs/comm*

Johnson PA

Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384-92

Jones JB

Meta-analyses: caveat lector, 15(2):61-2 *abs/comm*

Julien S

Diagnosis and management of preeclampsia, 15(12):607-14

Kallwitz ER

Management of the cirrhotic patient prior to liver transplantation, 15(8):393-400

Kamat PP

Barrett's esophagus: evaluation and management, 15(8):402-11

Kanwar M

Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20

Kaplowitz PB

Length of stay safely reduced with a diabetes clinical pathway in an urban children's hospital, 15(8):384-92

Katz JT

A program for reducing noneducational resident activities at an academic medical center, 15(1):25-30

Keyhani S

Another reason to drink your morning cup of coffee, 15(9):429-30 *abs/comm*

Do Medicaid enrollees in managed care plans receive similar care compared with commercial populations?, 15(1):8-9 *abs/comm*

FDA drug review deadlines: a safety concern?, 15(5):224-5 *abs/comm*

Kirch MA

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92

Kirchhoff AC

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Kirk JK

Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6

Klem ML

Optimizing medication adherence in older patients: a systematic review, 15(12):595-606

Koss RG

From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37

Kravet S

Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9

Kumar CB

Intensive intervention reduces mortality in type 2 diabetes, 15(4):171-3 *abs/comm*

Ladson D

Generalized anxiety disorder in adults, 15(7):353-63

Laing M

A program for reducing noneducational resident activities at an academic medical center, 15(1):25-30

Lamb P

Optimizing telemetry utilization in an academic medical center, 15(9):435-40

Laramie D

Periprocedural anticoagulation management in orthopedic patients: overview and description of a program utilizing pharmacogenetics, 15(4):183-90

Lee JC

Optimizing telemetry utilization in an academic medical center, 15(9):435-40

Levie J

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Lippold S

Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Litvin C

To err is human continued: a failure of follow-up, 15(1):21-3

Loeb JM

From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37

Malaty HM

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92

Manley M

Clinical system elements and patient reports of tobacco cessation counseling, 15(10):485-92

Martin KD

Isopropyl alcohol left on the skin falsely lowers capillary glucose values, 15(4):179-81

McCotter P

Supporting health care workers after medical error: considerations for health care leaders, 15(5):240-7

McKinley L

Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71-85

McMahon GT

Cardioprotective strategies for patients with type 2 diabetes, 15(3):133-46

Megginsong GC

Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43

Milsom VA

Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536-44

Minnick S

Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20

Montgomery D

Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34

Morton DJ

From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37

Mui C

New and emerging treatments in smoking cessation, 15(11):545-52

The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470-1 *abs/comm*

Murff HJ

Combination therapies are more effective than monotherapy for mild to moderate acne, 15(1):9-10 *abs/comm*

Musante D

Anorexia nervosa: role of the primary care physician, 15(9):450-60

Nackers LM

Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536-44

Naglak M

Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293-6

Newton RL Jr

Supporting behavior change in overweight patients: a guide for the primary care physician, 15(11):536-44

O'Malley K

Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92

Organist LA

Optimizing medication adherence in older patients: a systematic review, 15(12):595-606

Pantoja PM

Patient-centered care and outcomes after myocardial infarction, 15(3):119-24

Parks CM

Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43

Pastor W

Length of stay safely reduced with a diabetes

- clinical pathway in an urban children's hospital, 15(8):384-92
- Patterson B**
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34
- Pratt DS**
Evaluation and management of hepatitis B virus infection, 15(3):147-55
- Rachelefsky L**
A program for reducing noneducational resident activities at an academic medical center, 15(1):25-30
- Raji A**
Early intervention to prevent type 2 diabetes, 15(1):39-47
- Rand E**
Optimizing telemetry utilization in an academic medical center, 15(9):435-40
- Reed MC**
Coronary revascularization in patients with stable or asymptomatic coronary artery disease: a review of the COURAGE trial, 15(7):344-52
- Rohrer JE**
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66-70
- Rosman HS**
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20
- Rubal BJ**
Optimizing telemetry utilization in an academic medical center, 15(9):435-40
- Rudolph JL**
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502-11
- Ryan C**
Optimizing telemetry utilization in an academic medical center, 15(9):435-40
- Safdar N**
Measures to prevent ventilator-associated pneumonia: which are efficacious?, 15(2):71-85
- Saravolatz L**
Inpatient cardiac telemetry monitoring: are we overdoing it?, 15(1):16-20
- Schaefer CT**
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31
- Schafer AL**
Interventions to improve treatment of osteoporosis following fracture, 15(12):587-94
- Schlenk EA**
Optimizing medication adherence in older patients: a systematic review, 15(12):595-606
- Schlichting JA**
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31
- Schreiber K**
Postoperative delirium: overview and opportunities to optimize outcomes, 15(10):502-11
- Sekhobo JP**
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494-501
- Sellmeyer DE**
Interventions to improve treatment of osteoporosis following fracture, 15(12):587-94
- Setodji C**
Patient-centered care and outcomes after myocardial infarction, 15(3):119-24
- Shah NR**
Diagnosis of advanced-stage cancer more likely in the uninsured and Medicaid-insured, 15(6):272-4 *abs/comm*
- Intensive intervention reduces mortality in type 2 diabetes, 15(4):171-3 *abs/comm*
- Meta-analyses: caveat lector, 15(2):61-2 *abs/comm*
- The POISE trial: casting more doubt on the benefits of perioperative beta blockers, 15(10):470-1 *abs/comm*
- Sherman SE**
New and emerging treatments in smoking cessation, 15(11):545-52
- Simpson AG**
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43
- Siraj ES**
A young woman with a thyroid nodule, 15(9):443-9
- Update on the diagnosis and treatment of hyperthyroidism, 15(6):298-307
- Smith A**
Evaluation of quality of diabetes care in a multiethnic, low-income population, 15(6):281-6
- Smith DG**
Use of a wireless nurse alert fall monitor to prevent inpatient falls, 15(6):293-6
- Sun BC**
Patient-centered care and outcomes after myocardial infarction, 15(3):119-24
- Swanson LM**
Helping patients with insomnia: a guide for the primary care physician, 15(4):191-203
- Tenner CT**
To err is human continued: a failure of follow-up, 15(1):21-3
- Tkachenko M**
Recurrent abdominal pain symptoms in childhood: a population-based study, 15(6):287-92
- Trikudanathan S**
Cardioprotective strategies for patients with type 2 diabetes, 15(3):133-46
- Triola MM**
Achievement of blood lipid targets in patients informed of their coronary risk, 15(3):115-6 *abs/comm*
- Chronic use of thiazolidinediones increases fracture risk, 15(7):322-3 *abs/comm*
- Drug-eluting stents improve long-term nonfatal outcomes, 15(11):526, 528 *abs/comm*
- Veloski C**
A young woman with a thyroid nodule, 15(9):443-9
- Viette CR**
Implementation of an electronic health record with an embedded quality improvement program to improve the longitudinal care of outpatients with coronary artery disease, 15(7):338-43
- Wade T**
Change in body mass index over 1 year in a family medicine clinic: a retrospective analysis of medical records, 15(2):66-70
- Walter EB**
Improving influenza vaccine coverage among asthmatics: a practice-based research network study, 15(5):227-34
- Wang C**
Evaluation of a diabetes case management intervention in an underserved population: a retrospective cohort study at a health disparities collaborative site, 15(10):494-501
- Waterman AD**
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240-7
- White AA**
Supporting health care workers after medical error: considerations for health care leaders, 15(5):240-7
- Williams SC**
From bad to better: a qualitative assessment of low-performing hospitals that improved their smoking cessation counseling performance, 15(7):329-37
- Wilson J**
Biologic agents in the treatment of Crohn's disease, 15(1):31-8
- Wright S**
Facilitators and barriers involved in conducting hospital-based research across multiple sites, 15(5):235-9
- You WB**
Diagnosis and management of preeclampsia, 15(12):607-14
- Youssef NN**
Functional abdominal pain in children, 15(5):248-58
- Zaman**
Primary care management of the patient with cirrhosis, 15(11):554-63
- Zhang JX**
Hypertension and hyperlipidemia management in patients treated at community health centers, 15(3):125-31

Copyright 2008 by Turner White Communications Inc., Wayne, PA. All rights reserved.