

Chronic Kidney Disease and Coronary Heart Disease

To the Editor:

We read with interest the review of chronic kidney disease (CKD) by Balogun and Bolton [1] in the December 2004 issue of JCOM. We think that the importance of CKD as a major risk factor for coronary heart disease should be emphasized. The high rate of cardiovascular-related morbidity and mortality in patients with CKD not yet requiring dialysis is well established [2]. Many risk factors such as diabetes, hypertension, hyperlipidemia, and advanced age are present in patients with CKD, but CKD is an independent risk factor. Manjunath et al showed that each 10 mL/min per 1.73 m² decrease in the glomerular filtration rate correlates independently with a 5% higher cardiovascular risk [3]. In patients with known risk factors for coronary heart disease, even mild renal impairment is independently associated with development of cardiovascular disease [4,5]. CKD may be an even greater risk factor for coronary heart disease than diabetes mellitus. In the Bypass Angioplasty Revascularization Investigation (BARI) trial, after 7 years of follow-up of patients who underwent revascularization, 85% of 611 patients with diabetes were free from cardiac death, but only 77% of 46 patients with CKD and no diabetes mellitus were free from cardiac death [6]. The Tehran Lipid and Glucose Study showed that although CKD patients had a lower prevalence of major coronary artery disease risk factors compared with diabetes patients, patients with moderate CKD had a higher prevalence of electrocardiogram-defined coronary artery disease: 19.7% versus 14.7% in patients with diabetes [7]. The management of patients with CKD should include an intense focus on coronary heart disease risk factor modification.

Arash Rashidi, MD

*Department of Internal Medicine
Fairview Hospital
Cleveland Clinic Health System
Cleveland, OH*

Dale S. Adler, MD

*Division of Cardiology
Department of Internal Medicine
University Hospitals of Cleveland
Case Western Reserve University, School of Medicine
Cleveland, OH*

References

- Balogun RA, Bolton WK. Chronic kidney disease: a brief review for the primary care physician. *J Clin Outcomes Manage* 2004;11(12):784-90.
- Shlipak MG, Fried LF, Crump C, et al. Cardiovascular disease risk status in elderly persons with renal insufficiency. *Kidney Int* 2002;62:997-1004.
- Manjunath G, Tighiouart H, Coresh J, et al. Level of kidney function as a risk factor for cardiovascular outcomes in the elderly. *Kidney Int* 2003;63:1121-9.
- Pinkau T, Hilgers KF, Veelken R, Mann JF. How does minor renal dysfunction influence cardiovascular risk and the management of cardiovascular disease? *J Am Soc Nephrol* 2004; 15:517-23.
- Culleton BF, Larson MG, Wilson PW, et al. Cardiovascular disease and mortality in a community-based cohort with mild renal insufficiency. *Kidney Int* 1999;56:2214-9.
- Szczzech LA, Best PJ, Crowley E, et al. Outcomes of patients with chronic renal insufficiency in the bypass angioplasty revascularization investigation. *Circulation* 2002;105:2253-8.
- Rashidi A, Adler DS, Ghanbarian A, Aziz F. Is chronic kidney disease a cardiovascular risk equivalent, as diabetes? Data from Tehran Lipid and Glucose Study (TLGS). In press 2005.

In reply:

We agree with Drs. Rashidi and Adler that CKD is a major risk factor for progression of CHD as well as other cardiovascular disease. It has been widely accepted that cardiac-related events are the number one cause of death in CKD stage 5 patients treated with hemodialysis. More recent data suggest that earlier-stage CKD can be regarded as a nontraditional and independent risk factor for mortality from CHD. Muntner et al reporting on the Second National Health and Nutrition Examination Survey Mortality Study showed that in a typical U.S. population, there was an independent association between CKD and increased all-cause mortality related to cardiac disease [1]. Go et al also showed an association between earlier stages of CKD and risks of cardiovascular events, hospitalization, and mortality [2]. Other recent articles have also demonstrated the association between CKD and cardiovascular disease [3-6]. We agree with Drs. Rashidi and Adler that the management of CKD should include management of comorbid conditions with emphasis on cardiovascular risk reduction through blood pressure control (to a goal of 130/80 mm Hg), lifestyle and dietary changes, smoking cessation, and so on. The clinical practice guidelines published by the Renal Physicians Association and the National Kidney Foundation provide evidence-based recommendations for this specialized group of patients.

Rasheed A. Balogun, MD

W. Kline Bolton, MD

*Division of Nephrology
University of Virginia Health System
Charlottesville, VA*

References

- Muntner P, He J, Hamm L, et al. Renal insufficiency and subsequent death resulting from cardiovascular disease in the United States. *J Am Soc Nephrol* 2002;13:745-53.
- Go AS, Chertow GM, Fan D, et al. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 2004;351:1296-305.
- Muntner P, He J, Astor BC, et al. Traditional and nontraditional

risk factors predict coronary heart disease in chronic kidney disease: results from the Atherosclerosis Risk in Communities study. *J Am Soc Nephrol* 2005;16:529–38.

4. Weiner DE, Tighiouart H, Stark PC, et al. Kidney disease as a risk factor for recurrent cardiovascular disease and mortality. *Am J Kidney Dis* 2004;44:198–206.
5. Weiner DE, Tighiouart H, Amin MG, et al. Chronic kidney disease as a risk factor for cardiovascular disease and all-cause mortality: a pooled analysis of community-based studies. *J Am Soc Nephrol* 2004;15:1307–15.
6. Foley RN, Murray AM, Li S, et al. Chronic kidney disease and the risk for cardiovascular disease, renal replacement, and death in the United States medicare population, 1998 to 1999. *J Am Soc Nephrol* 2005;16:489–95.

Copyright 2005 by Turner White Communications Inc., Wayne, PA. All rights reserved.