

# HOSPITAL PHYSICIAN®

## INFECTIOUS DISEASES BOARD REVIEW MANUAL

### STATEMENT OF EDITORIAL PURPOSE

The *Hospital Physician Infectious Diseases Board Review Manual* is a study guide for fellows and practicing physicians preparing for board examinations in infectious diseases. Each manual reviews a topic essential to current practice in the subspecialty of infectious diseases.

### PUBLISHING STAFF

#### PRESIDENT, GROUP PUBLISHER

Bruce M. White

#### EDITORIAL DIRECTOR

Debra Dreger

#### SENIOR EDITOR

Bobbie Lewis

#### EDITOR

Tricia Faggioli, ELS

#### ASSISTANT EDITOR

Farrowh Charles

#### EXECUTIVE VICE PRESIDENT

Barbara T. White

#### EXECUTIVE DIRECTOR OF OPERATIONS

Jean M. Gaul

#### PRODUCTION DIRECTOR

Suzanne S. Banish

#### PRODUCTION ASSISTANT

Nadja V. Frist

#### ADVERTISING/PROJECT DIRECTOR

Patricia Payne Castle

#### SALES & MARKETING MANAGER

Deborah D. Chavis

#### NOTE FROM THE PUBLISHER:

This publication has been developed without involvement of or review by the American Board of Internal Medicine.

## Common Clinical Problems in Returned Travelers

### Series Editor:

**Mary E. Klotman, MD**

*Chief, Division of Infectious Diseases, Mt. Sinai School of Medicine, New York, NY*

### Contributor:

**Daniel Caplivski, MD**

*Director, Travel Medicine Program, Assistant Professor, Division of Infectious Diseases, Mount Sinai School of Medicine, New York, NY*

## Table of Contents

Introduction .....	2
Fever .....	2
Skin Lesions .....	5
Gastrointestinal Infections .....	9
Conclusion .....	11
References .....	11

Cover Illustration by Kathryn K. Johnson

Copyright 2008, Turner White Communications, Inc., Strafford Avenue, Suite 220, Wayne, PA 19087-3391, www.turner-white.com. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, mechanical, electronic, photocopying, recording, or otherwise, without the prior written permission of Turner White Communications. The preparation and distribution of this publication are supported by sponsorship subject to written agreements that stipulate and ensure the editorial independence of Turner White Communications. Turner White Communications retains full control over the design and production of all published materials, including selection of topics and preparation of editorial content. The authors are solely responsible for substantive content. Statements expressed reflect the views of the authors and not necessarily the opinions or policies of Turner White Communications. Turner White Communications accepts no responsibility for statements made by authors and will not be liable for any errors of omission or inaccuracies. Information contained within this publication should not be used as a substitute for clinical judgment.

# Common Clinical Problems in Returned Travelers

Daniel Caplivski, MD

## INTRODUCTION

The increasing frequency of international travel has led to an increased number of patients presenting with illnesses acquired abroad.<sup>1</sup> It is therefore important that practicing physicians have an understanding of the common problems facing returned travelers. This review will discuss some of the principles that guide evaluation and management of these patients in a format that is based on actual cases from the Mount Sinai School of Medicine Travel Medicine Program.

## FEVER

Fever can be an important sign in several life-threatening infections. The question “Could this be malaria?” must always be posed in the setting of the febrile returned traveler because of the high mortality associated with untreated malaria due to *Plasmodium falciparum*. The absence of fever should not discount the possibility of malaria infection, as some patients may present with more subtle symptoms.<sup>1</sup>

When considering the possibility of travel-related illness, one must entertain the possibility that more local causes of infection are in fact the true culprit. A careful history of the patient’s itinerary, activities, and exposures as well as the chronicity of the symptoms will often help to differentiate among the various causes of fever without localizing symptoms.<sup>1</sup>

## CASE 1



A 43-year-old man from Nigeria returned to his home country for a 6-month stay while working for the United Nations. He had received appropriate pretravel vaccinations but was not taking malaria prophylaxis while there. Most of his daily activities were confined to offices, but he had made several trips to smaller rural villages in which he had spent a large portion of his time outdoors. The night before his return, he developed fever and headache. His symptoms then

progressed to include chills, myalgias, and extreme fatigue. By the time he presented to the emergency department, he had also noted a darker color to his urine and a significant icterus (**Figure 1**). His laboratory studies revealed several abnormalities (abnormal results in italics):

- White blood cell count: *11,300/μL* (71% neutrophils, *12% bands*)
- Hemoglobin: *10.8 g/dL*
- Hematocrit: *30%*
- Platelet count: *29,000/μL*
- Sodium: *127 mEq/L*
- Potassium: *4.2 mEq/L*
- Chloride: *91 mEq/L*
- Carbon dioxide, total: *18 mEq/L*
- Blood urea nitrogen: *84 mg/dL*
- Creatinine: *5.7 mg/dL*
- Lactate dehydrogenase: *1757 U/L*
- Alanine aminotransferase: *59 U/L*
- Aspartate aminotransferase: *131 U/L*
- Bilirubin, total: *10.5 mg/dL*
- Bilirubin, direct: *5.4 mg/dL*
- Haptoglobin: *10 mg/dL*

On peripheral blood smear, there were multiple ring forms with a 20% parasitemia (**Figure 2**).

- **What is the diagnosis? What is the most likely species?**

This clinical presentation is consistent with severe malaria from *Plasmodium falciparum*. On the peripheral blood smear, the infected cells are the same size as the noninfected cells and there is an example of a cell infected by more than 1 *Plasmodium* parasite. These findings together with the clinical picture of a patient with severe malaria allow us to distinguish infection with *P. falciparum* from infection with other species of malaria. On peripheral blood smear, both *P. vivax* and *P. ovale* will infect immature erythrocytes that have not fully developed their cytoskeleton. These infected cells will appear larger than the noninfected adjacent cells. In *P. falciparum*, the schizont phase is rarely seen on